

Asthma

A study to compare different doses of MSTT1041A with a “placebo” – in patients with severe asthma

A Study to Assess the Efficacy and Safety of MSTT1041A in Participants With Uncontrolled Severe Asthma

Trial Status
Completed

Trial Runs In
15 Countries

Trial Identifier
NCT02918019 2016-001549-13
GB39242

The source of the below information is the publicly available website [ClinicalTrials.gov](https://clinicaltrials.gov). It has been summarised and edited into simpler language.

Trial Summary:

This is a Phase IIb, randomized, placebo-controlled, double-blind, multicenter, multi-arm study which will evaluate efficacy, safety, and pharmacokinetic of MSTT1041A compared with placebo as add-on therapy in participants with severe, uncontrolled asthma who are receiving medium- or high-dose inhaled corticosteroid (ICS) therapy and at least one of the following additional controller medications: long-acting beta-agonists (LABA), leukotriene modifier (LTM), long-acting muscarinic antagonist (LAMA), or long-acting theophylline preparation. The total duration of this study for each participant is approximately 70 weeks including screening, run-in, treatment, and follow-up.

Hoffmann-La Roche
Sponsor

Phase 2
Phase

NCT02918019 2016-001549-13 GB39242
Trial Identifiers

Eligibility Criteria:

Gender
All

Age
≥18 Years & ≤ 75 Years

Healthy Volunteers
No

This clinical trial was done to study a new medicine called, “MSTT1041A”, for the treatment of patients with severe asthma. This study was done to find the dose of MSTT1041A that was effective for reducing the number of asthma exacerbations. Other questions that researchers had included finding the dose of MSTT1041A that was effective for improving other symptoms for asthma patients; time it took for MSTT1041A to be distributed in the body; whether MSTT1041A was safe for asthma patients; and whether

ForPatients

by Roche

MSTT1041A caused the immune system to make antibodies against this medicine. There were 502 patients with asthma who took part in this study at 182 study centers in 15 countries.